Draeger (Models B, C, and GS) Anesthesia Machine Checklist Northwestern Memorial Hospital, Chicago, IL

- 1. ADJUST FLOW ON SCAVENGER!
- 2. Check Ambu bag (Including backup Oxygen Source)
- 3. Fill vaporizers.
- 4. Check CO₂ Absorbant (depletion and dessication).
- 5. Remove oxygen analyzer fuel cell from breathing circuit.
- 6. Turn on anesthesia machine and monitors.
- 7. Calibrate O₂ analyzer to 21% and reinstall fuel cell.
 - a. Confirm Low O₂ Alarm Function and Low Limit (30%)
- 8. Disconnect central gas supply hose for O₂.
- 9. Check pressure in O₂ cylinder(s) (>1000 psi).
- 10. Bleed O₂ pressure so tank pressure gauge reads zero.
- 11. Connect central gas supply hose for O₂.
 - a. Confirm central supply gas pressures of ±50 psi
 - b. Confirm full 10 lpm for O₂, N₂O, and air (Helium if so equipped)
 - i. Turn on N₂O first to confirm zero flow

12. Check ventilator function:

- a. Connect reservoir bag to Y-piece of breathing circuit
- b. Position selector valve to ventilator (auto)
- c. Turn on ventilator; set at an appropriate:

Tidal Volume, Rate, I:E ratio and Pressure Limit

- d. Check unidirectional valve function
- e. Remove/Replace reservoir bag and occlude outlet at Y-piece
- f. Fill bellows and perform high pressure check
- a. Turn off ventilator
- h. Confirm Ventilator Disconnect Alarm

13. Leak test the breathing circuit:

- a. Position selector valve to bag
- b. Close APL (adjustable pressure limit) valve
- c. Confirm zero flow on flowmeters
- d. Occlude outlet at Y-piece
- e. Pressurize circuit to 30 cmH₂O using O₂ flush valve
- f. Turn on and off each vaporizer Confirming sustained circuit pressure of 10 seconds for each vaporizer
- g. Open APL valve to slowly decompress circuit
- 14. Check APL valve and scavenger:
 - a. With APL valve open and Y-connector occluded press O₂ flush valve and confirm pressure of < 14 cmH₂O

15. Check suction:

- a. Check connections on suction cannister
- b. Check adequacy of vacuum
- c. Connect Yankauer suction tip
- 16. Check laryngoscope blade.
- 17. Prepare endotracheal tube.
- 18. Check drugs and syringes (keep empty vials and ampules).
- 19. Document completed checkout on anesthesia record.
- 20. ANESTHESIA TIME-OUT.

Northwestern University, Chicago Tom C. Kreicie, MD Sample

Revised: September 1, 2007