

# ANESTHESIOLOGY



## Preoperative Cognitive Dysfunction

**\* It's More Common Than You Think**

**Cognitive impairment increases with age...**

Age Group	Cognitive Impairment (%)
70-79 yr	5%
80-89 yr	24%
90+ yr	40%

**...and preoperative cognitive dysfunction is associated with adverse outcomes.**

1 in 3

surgical patients are 65 or older

**Although many patients are at risk, the impact of cognitive impairment on elective surgery outcomes is unknown.**

**Culley *et al.*<sup>1</sup> screened orthopedic patients over 65 for probable cognitive impairment using the MiniCog.**

+

**2 points** for drawing clock

**1 point** per word recalled

= MiniCog Score

0-2: Positive screen  
3-5: Negative screen

**24% of patients assessed had probable cognitive impairment.**

**Probable cognitive impairment was associated with worse surgical outcomes:**

**67%**  
(vs. 34% without impairment)

**Discharge other than home**

**21%**  
(vs. 7% without impairment)

**Postoperative delirium**

**1.12**  
days  
(95% CI, 0.67-1.58)

**Longer hospital stay**

**These data support screening at-risk patients to identify those at high risk for adverse outcomes. Additional work is needed to define strategies to mitigate these risks.**

Infographic created by Jonathan P. Wanderer, Vanderbilt University Medical Center, and James P. Rathmell, Brigham and Women's Health Care/Harvard Medical School. Illustration by Annemarie Johnson, Vivo Visuals. Address correspondence to Dr. Wanderer: jonathan.p.wanderer@vanderbilt.edu.

1. Culley DJ, Flaherty D, Fahey MC, Rudolph JL, Javedan H, Huang C-C, Wright J, Bader AM, Hyman BT, Blacker D, Crosby G: Poor performance on a preoperative cognitive screening test predicts postoperative complications in older orthopedic surgical patients. *ANESTHESIOLOGY* 2017; 127:765-74